

EXPRESS MAIL CERTIFICATE

Date 3/7/01 Label No. EL706723006 #18

5 I hereby certify that, on the date indicated above I deposited
this paper or fee with the U.S. Postal Service & that it was
addressed for delivery to the Assistant Commissioner of
Patents & Trademarks, Washington, DC 20231 by
"Express Mail Post Office to Addressee" service.



10 J. DiLullo J. DiLullo
Name (Print) Signature

07278
PATENT TRADEMARK OFFICE

15

2427/1G685US1

20 **HIGHLY EFFICIENT GENE TRANSFER INTO HUMAN REPOPULATING
STEM CELLS BY *RD114* PSEUDOTYPED RETROVIRAL VECTOR PARTICLES**

This application claims priority under 35 U.S.C. §119(e) of provisional
patent application Serial NO. 60/187,534, filed March 7, 2000, which is incorporated
herein by reference in its entirety.

25 The research leading to the present invention was supported in part by
National Institutes of Health Grant P01 HL 53749. The United States Government may
have certain rights in the invention.

FIELD OF THE INVENTION

30 The present invention relates to a method for efficiently introducing
exogenous genes into stem cells, particularly human repopulating stem cells, using
retroviral vector particles pseudotyped with the feline endogenous retrovirus (RD114)
envelope protein. The gene transfer method of the present invention is important for
somatic cell gene therapy, for studying the differentiation of various cell lineages, and for
35 creating animal models of various human stem cell conditions.

BACKGROUND OF THE INVENTION

Use of Retroviral Vectors for Gene Therapy

Gene therapy is a novel method under investigation for the treatment of

genetic, metabolic and neurologic diseases, cancer and AIDS. The primary goal of gene therapy is to deliver a specific gene to a pre-determined target cell, and to direct expression of such a gene in a manner which will result in a therapeutic effect.

Compared to more traditional methods of gene transfer, retroviral vectors
 5 are extraordinarily efficient gene delivery vehicles (for reviews *see* Vile *et al.*, Br. Med. Bull. 1995, 51:12-30; Klimatcheva *et al.*, Front Biosci., 1999, 4:D481-96). The retroviral genome becomes integrated into host chromosomal DNA, ensuring its long-term persistence and stable transmission to all future progeny of the transduced cell and making retroviral vector suitable for permanent genetic modification. Up to 8 kilobases of foreign
 10 gene sequence can be packaged in a retroviral vector particles, which is more than enough for most gene therapy applications. The ability of retroviral vector particles to cause no detectable harm while entering their target cells represents another therapeutically important property. Indeed, there have been no reported short- or long-term toxicity problems associated with the use of the retroviral vectors in human gene therapy trials,
 15 now dating back to 1989. In addition, these vectors can be manufactured in large quantities, which allows their standardization and use in pharmaceutical preparations.

In view of the properties described above, it is not surprising that retroviral vectors have been selected as the vectors of choice in about 80% of the clinical gene therapy trials that have been approved to date. The most successful of these clinical
 20 experiments have used T cell-directed retroviral vector-mediated gene transfer to cure a severe combined immunodeficiency (SCID) caused by adenosine deaminase (ADA) deficiency (genetic disease) in humans (for review *see* Onodera *et al.*, Acta Haematol., 1999, 101:89-96).

Most of the commonly used retroviral vectors are of oncoviral origin.
 25 These vectors require cell division in order to achieve genome integration and long-term gene expression. Accordingly, lentiviral vectors (*e.g.*, HIV-based), which are retroviruses capable of productively infecting non-dividing cells, have been suggested as an alternative approach to successful gene transfer in quiescent somatic cells (Miller *et al.*, 1990, Mol. Cell Biol., 10:4239-4242; Klimatcheva *et al.*, *supra*).

30 **Retrovirus vectors.** Introduction of genes into host animals is described in U.S. Patent No. 5,399,346; Mann *et al.*, Cell, 1983, 33:153; U.S. Patent Nos. 4,650,764 and 4,980,289; Markowitz *et al.*, J. Virol., 1988, 62:1120; U.S. Patent No. 5,124,263;

European Publication Nos. EP 453 242 and EP178 220; Bernstein *et al.*, Genet. Eng., 1985, 7:235; McCormick, BioTechnology, 1985, 3:689; PCT Publication No. WO 95/07358; and Kuo *et al.*, Blood, 1993, 82:845. Retroviral vectors can be constructed from different types of retrovirus, such as HIV (human immunodeficiency virus), MoMuLV ("murine Moloney leukemia virus"), MSV ("murine Moloney sarcoma virus"), HaSV ("Harvey sarcoma virus"); SNV ("spleen necrosis virus"); RSV ("Rous sarcoma virus"), and Friend virus. In the last decade, several retroviral vector systems, derived from chicken or murine retroviruses, have been developed for the expression of various genes (for reviews see Temin, 1987; Gilboa, 1990; Robbins and Ghivizzani, Pharmacol. Ther., 1998, 80:35-47).

10 The majority of existing gene therapy protocols use simple retroviral vectors based on Murine Leukemia Virus (MLV). These vectors are able to infect many different cell types, and their retroviral promoter, which is often used to control the expression of a heterologous (therapeutic) gene, is active in a wide range of different cell types.

Retroviral vectors can be constructed to function as infectious particles or

15 to undergo a single round of transfection. In the former case, the virus is modified to retain all of its genes except for those responsible for oncogenic transformation properties, and to express the heterologous gene. Non-infectious viral vectors are manipulated to destroy the viral packaging signal from the genes encoding viral structural proteins, retaining the structural genes required to package the co-introduced virus engineered to

20 contain the heterologous gene and the packaging signals. Thus, the viral particles that are produced are not capable of producing additional virus.

Packaging cell lines contain the genes for retrovirus replication and assembly, *e.g.*, gag, pol, and/or env. When transfected with a retroviral genome containing a gene of interest and assembly signals, the packaging cells become producer

25 cells. Suitable packaging cell lines have been described in the prior art, in particular the cell line PA317 (U.S. Patent No. 4,861,719); the PsiCRIP cell line (PCT Publication No. WO 90/02806) and the GP+envAm-12 cell line (PCT Publication No. WO 89/07150). In addition, the recombinant retroviral vectors can contain modifications within the LTRs for suppressing transcriptional activity as well as extensive encapsidation sequences which

30 may include a part of the gag gene (Bender *et al.*, J. Virol., 1987, 61:1639). Recombinant retroviral vectors are purified by standard techniques known to those having ordinary skill in the art.

Lentivirus vectors. Lentiviral vectors, which are a particular type of retrovirus vector, have also been used as agents for the direct delivery and sustained expression of a transgene in several tissue types, including brain, retina, muscle, liver, and blood. The vectors can efficiently transduce dividing and nondividing cells in these tissues, and maintain long-term expression of the gene of interest. For a review, *see*, Naldini (Curr. Opin. Biotechnol., 1998, 9:457-63; *see also* Zufferey, *et al.*, J. Virol., 1998, 72:9873-80). Lentiviral packaging cell lines are available and known generally in the art. They facilitate the production of high-titer lentivirus vectors for gene therapy. An example is a tetracycline-inducible VSV-G pseudotyped lentivirus packaging cell line that can generate virus particles at titers greater than 10^6 IU/ml for at least 3 to 4 days (Kafri, *et al.*, J. Virol., 1999, 73: 576-584). The vector produced by the inducible cell line can be concentrated as needed for efficiently transducing non-dividing cells *in vitro* and *in vivo*.

However, despite this impressive record, there is still a great need for the development of new, improved retroviral vectors and packaging systems to fuel further advances in the field of human gene therapy.

Structure of a Retroviral Vector

Retroviral vectors are infection-competent viral particles that contain functional packaging signals and a crippled genome in which most or all of the retroviral protein coding sequences have been replaced with the gene(s) of interest. As a result, upon infection of a target cell, such viruses cannot undergo more than one round of replication in the absence of a helper virus. Retroviral vector particles are produced by helper cells (also called producer cells), which contain constructs expressing all retroviral proteins necessary for particle production and replication (*i.e.*, at least three proteins: gag, pol, and env). After the introduction (transfection) of the RNA genome of a retroviral vector into such helper cells, it becomes encapsulated into virus particles (due to the presence of specific packaging signals in its RNA) having a complete set of retroviral proteins sufficient to support infection and a single round of replication and carrying a genome containing gene(s) of interest, and the packaging signals required to incorporate the genome into the viral particle. Upon their release from the helper cells, the retroviral vector particles can be isolated from the culture medium and used to infect (transduce) various types of target cells. Following transduction, the RNA genome is reverse

transcribed into DNA and the DNA copy (provirus) is integrated into the host genome.

Safe and efficient retroviral-mediated gene transfer systems that are suitable for somatic gene therapy are likely to involve the development of alternative, cell type-targeted vectors containing the cell type-specific retroviral envelope protein (*env*), which, together with the availability of its cognate receptor, determines infectivity. Cell type targeting may be also achieved by using cell-specific promoters to limit the induction of expression of retroviral vector-encoded genes (for review *see* Gunzburg *et al.*, Cytokines Mol. Ther., 1996, 2:177-84).

10 Use of Stem Cells for Retrovirus-Mediated Gene Delivery

The potential use of gene-modified repopulating stem cells as vehicles for retrovirus-mediated gene delivery was first illustrated by the demonstration that hematopoietic systems of lethally irradiated mice can be reconstituted with retroviral vector-transduced syngenic bone marrow cells, and that these cells can, in turn, provide 15 genetically marked progeny, which persist in blood and marrow over extended periods of time (Chu *et al.*, J. Mol. Med., 1998, 76:184-92; Medin and Karlsson, Immunotechnology, 1997, 3:3-19).

The ability to transfer genes into stem cells of hematopoietic, neural or muscular origin would provide an opportunity for somatic gene therapy of various 20 malignant and non-malignant disorders. Thus, gene transfer to primary hematopoietic cells may be used to treat diseases that affect bone marrow and peripheral blood function (for reviews *see* Anderson, Science, 1984, 226:401-409; Sorrentino *et al.*, pp. 351-426, *In: The Development of Gene Therapy*, T. Friedmann ed., New York: Cold Spring Harbor Laboratory Press, 1999). Examples of genetic diseases which are potentially curable by 25 transgene-mediated delivery of a normal gene product include: X-linked agammaglobulinemia (Vetrie *et al.*, Nature, 1993, 361:226; Tusukada *et al.*, Cell, 1993, 72:279), ADA deficiency (Anderson, Science, 1992, 256:808), hemophilia (factor VIII and factor IX deficiency) (Miller, Blood, 1990, 76:271; Hoeben *et al.*, Thromb. Haemost., 1992, 67:341; Hoeben *et al.*, Hum. Gene Ther., 1993, 4:179; Herzog and High, Curr. Opin. Hematol., 1998, 5:321-6), and lipoprotein lipase deficiency (Hayden and Ma, Mol. Cell. Biochem., 1992, 113:171; Auwerx *et al.*, Crit. Rev. Clin. Lab. Sci., 1992, 29:243). Indeed, 30 transfer of the gene coding for glucocerebrosidase (GC) via a retroviral vector (MFG-GC)

to human CD34 cells obtained from patients with Gaucher's disease was shown to result in correction of the enzyme deficiency in these cells (Barranger *et al.*, Baillieres Clin. Haematol., 1997, 10:765-78). Similarly, hematopoietic stem cells are an attractive target for gene therapy of AIDS because of their ability to generate a broad repertoire of mature T lymphocytes, as well as the monocytic cells (macrophages, dendritic cells and microglia), which are also involved in HIV-1 pathogenesis. A number of synthetic "anti-HIV-1 genes" have been developed which inhibit HIV-1 replication (Engel and Kohn, Front. Biosci., 1999, 4:e26-33). Gene transfer approaches can be also used in diagnosis and treatment of cancer, *e.g.*, to "mark" cancer cells to monitor their persistence *in vivo* in patients, to protect normal cells from toxic chemotherapeutic agents, to correct a genetic defect in or to confer a novel function on the cancer cell (Clay *et al.*, Pathol. Oncol. Res., 1999, 5:3-15). For example, preliminary experiments in transgenic mice have demonstrated that overexpression of the human multidrug resistance 1 (MDR1) gene results in a complete protection against high doses of cytostatic drugs (Fruehauf *et al.*, Recent Results Cancer Res., 1998, 144:93-115).

Several gene therapy protocols involving retroviral-vector-mediated transfer of exogenous genes into cells of hematopoietic lineage are in clinical trials. These include the introduction of the adenosine deaminase (ADA) gene into peripheral blood T cells of ADA deficient, severe combined immunodeficiency (SCID) patients (Sprent, Cell. Immunol., 1983, 7:10; Stevens *et al.*, J. Immunol., 1982, 128:844), and the introduction of marker genes or lymphokine genes into tumor infiltrating (TIL) T cells (Kasis *et al.*, Proc. Natl. Acad. Sci. USA, 1990, 87:473; Culver *et al.*, 1991, *ibid.*, 88:3155).

Existing Problems in Retroviral-Mediated Gene Transfer to Stem Cells

Extensive efforts have been invested in adapting oncoretroviral vectors for gene transfer into stem cells (Allay *et al.*, Nat. Med., 1998, 4:1136-1143; Bunting *et al.*, Nat. Med., 1998, 4:58-64; Persons *et al.*, Nat. Med., 1998, 4:1201-1205; Donahue *et al.*, Blood, 2000, 95:445-452; Dunbar *et al.*, Blood, 1995, 85:3048-3057; Kiem *et al.*, Blood, 1998, 92:1878-1886). Despite considerable success in murine models, retroviral-mediated gene transfer into human stem cells has been difficult to achieve due to low abundance of viral receptors, inhibition of retroviral transgene expression, and the relative quiescence of the target cell population (Kurre *et al.*, J. Virol., 1990, 73:495-500; Miller *et al.*, Mol. Cell

Biol., 1990, 10:4239-4242; Orlic *et al.*, Proc. Natl. Acad. Sci. USA, 1996, 93:11097-11102; Bhatia *et al.*, J. Exp. Med., 1997, 186:619-624; Glimm *et al.*, Blood, 1999, 94:2161-2168.; Hao *et al.*, Blood, 1996, 88:3306-3313). As explained above, the latter problem can be solved by the use of lentiviral vectors, which are able to infect non-

5 dividing cells. In addition, the recent evidence suggests that repopulating hematopoietic stem cells can be induced to divide in short-term cultures with high concentrations of cytokines (Bhatia *et al.*, *supra*; Glimm H, *et al.*, *supra*), thus opening the possibility to increase the efficiency of oncoviral-mediated transduction. However, upon stimulation these stem cells are likely to differentiate, thus defeating the goal of using them in the first

10 place.

It appears therefore that the retroviral receptor deficiency and the efficiency and persistence of retroviral transgene expression are the major remaining barriers to stem cell-targeted gene transfer.

Another factor which has complicated the development of stem cell-

15 targeted gene therapy approaches for human disorders is the lack of reliable *in vitro* surrogate assays for the repopulating cells. Thus, successful gene transfer into hematopoietic progenitors, as evaluated in *in vitro* clonogenic assays, has been a poor predictor of long-term gene transfer into repopulating stem cells in large animal models or human clinical trials (Dunbar *et al.*, *supra*). Cells capable of initiating long-term culture

20 (LTC-IC), particularly those which give rise to hematopoietic progenitors over 10-12 weeks, may be a much better surrogate for repopulating stem cells (Hao *et al.*, *supra*; Sutherland *et al.*, Proc. Natl. Acad. Sci. USA, 1990, 87:3584-3588). Another surrogate for stem cell targeted gene transfer are the primitive human hematopoietic cells that are able to establish hematopoiesis in immunodeficient mice, such as the non-obese diabetic/severe

25 combined immunodeficiency (NOD/SCID) mouse strain. Human NOD/SCID repopulating cells (SRC) become engrafted in various hematopoietic cell lineages and can be recovered from murine bone marrow months after transplantation (Kamel-Reid *et al.*, Science, 1988, 242:1706- 1709; Vormoor *et al.*, Blood, 1994, 83:2489-2497; Cashman *et al.*, Blood, 1997, 89:4307- 4316; Larochelle *et al.*, Nat. Med., 1996, 2:1329-1337). It

30 follows that gene transfer strategies resulting in retroviral marking of a significant proportion of the progeny of SRC would be a better predictor for stem cell gene transfer than surrogates evaluated by *in vitro* assays (Larochelle *et al.*, Nat. Med., 1996, 2:1329-

1337).

Most vector preparations utilized for gene transfer into human cells contain particles having amphotropic specificity based on the structure of their envelope protein. Amphotropic viral particles have a broad host range that includes human cells (Kurre *et al.*, J. Virol., 1999, 73:495-500). In contrast, ecotropic vector particles used for gene transfer into murine stem cells are unable to transduce human cells because of interspecies polymorphic variation in the cationic transporter which serves as the receptor for ecotropic vector particles (Albritton *et al.*, J. Virol., 1993, 67:2091-2096; Kizhatil *et al.*, J. Virol., 1997, 71:7145-7156). Despite their broad cell type specificity, amphotropic vector particles infect human hematopoietic stem cells very inefficiently. It has been recently demonstrated that amphotropic vector particles enter their target cells via interactions with a phosphate transporter (van Zeijl *et al.*, Proc. Natl. Acad. Sci. USA, 1994, 91:1168-1172; Miller *et al.*, Proc. Natl. Acad. Sci. USA, 1994, 91:78-82; Kavanaugh *et al.*, Proc. Natl. Acad. Sci. USA, 1994, 91:7071-7075), which is expressed at very low levels on primitive human hematopoietic cells (Orlic *et al.*, *supra*) and possibly other types of stem cells. To overcome this barrier to retroviral-mediated gene transfer, vector preparations pseudotyped with envelope proteins from other viruses have been generated and studied.

Oncoretroviral vectors pseudotyped with the envelope protein of the Gibbon Ape Leukemia Virus (GALV) were shown to transduce human clonogenic hematopoietic progenitor cells (Bauer *et al.*, Blood, 1995, 86:2379-2387) and NOD/SCID repopulating cells (SRC) from cord blood more efficiently than amphotropic vector particles (van Hennik *et al.*, Blood, 1998, 92:4013-4022; Marandin *et al.*, Hum. Gene Ther., 1998, 9:1497-1511; Conneally *et al.*, Blood, 1998, 91:3487-3493). In accord with these data, the GALV receptor, also a phosphate transporter, was found to be expressed at a somewhat higher level than the amphotropic receptor (Kiem *et al.*, Blood, 1998, 92:1878-1886; Bauer *et al.*, Blood, 1995, 86:2379-2387). Thus, in studies in the NOD/SCID model, approximately 25% of human CD45⁺ cells in multiple hematopoietic lineages of transplant recipients were positive for the vector genome as documented by PCR or transgene expression only after multiple exposures to viral vectors (van Hennik *et al.*, *supra*; Marandin *et al.*, *supra*; Schilz *et al.*, Blood, 1998, 92:3163-3171; Hennemann *et al.*, Experimental Hematology, 1999, 27:817-825). These data correlate with the results of larger animal studies in which GALV pseudotyped oncoretroviral vector particles show

higher frequency of gene transfer than amphotropic vector particles and are incorporated into 10% of myeloablated canine and baboon models (Kiem *et al.*, Blood, 1998, 92:1878-1886; Tisdale *et al.*, Blood, 1998, 92:1131-1141; Sellers *et al.*, Hum. Gene Ther., 1999, 10:633-640; Whitwam *et al.*, Blood, 1998, 92:1565-1575).

5 The G envelope protein of vesicular stomatitis virus (VSV-G), a rhabdovirus, has been used to pseudotype both lentiviral and oncoviral vector particles (Yang *et al.*, Hum. Gene Ther., 1995, 6:1203-1213; Emi *et al.*, J. Virol., 1991, 65:1202-1207; Friedmann *et al.*, Nat. Med., 1995, 1:275-277; Akkina *et al.*, J. Virol., 1996, 70:2581-2585; Naldini *et al.*, Science, 1996, 272:263-267). The VSV-G pseudotyped
10 particles can be concentrated 100-1,000 fold by centrifugation and they enter target cells via interaction with phospholipids which are found on all cell types (Burns *et al.*, Proc. Natl. Acad. Sci. USA, 1993, 90:8033-8037). It has been shown that both VSV-G pseudotyped oncoviral and lentiviral vector particles efficiently transduce hematopoietic stem cells (Evans *et al.*, Hum. Gene Ther., 1999, 10:1479-1489; Douglas *et al.*, Hum.
15 Gene Ther., 1999, 10:935-945; Case *et al.*, Proc. Natl. Acad. Sci. USA, 1999, 96: 2988-2993; Sinclair *et al.*, Gene Ther., 1997, 4:918-927). Successful gene transfer into SRC using VSV-G pseudotyped oncoretroviral or lentiviral vector particles has been also reported (Rebel *et al.*, Blood, 1999, 93:2217-2224; Miyoshi *et al.*, Science, 1999, 283:682-686). Despite the increased efficiency of gene transfer observed with VSV-G pseudotyped
20 particles compared with GALV pseudotyped and amphotropic vector particles, the use of these particles for therapeutic purposes is questionable. VSV-G protein is toxic to producer cells by causing the membrane fusion, and the inducible promoters must be used to control its expression (Yang *et al.*, *supra*; Ory *et al.*, Proc. Natl. Acad. Sci. USA, 1996, 93:11400-11406).

25 Vector particles pseudotyped with the envelope protein of the feline endogenous virus RD114 have been described (U.S. Patent No. 5,952,225). The RD114 retrovirus is a member of the large interference group 1 of retroviruses all of which use the same receptor on human cells (Sommerfelt *et al.*, Virology, 1990, 176:58-69) recently identified as a neutral amino acid transporter (Rasko *et al.*, Proc. Natl. Acad. Sci. USA,
30 1999, 96:2129-2134). In addition, murine RD114 pseudotyped vector particles have been shown transduce human hematopoietic cells at about the same efficiency as amphotropic vector, when the target cells were cultured with the producer cell line (Porter *et al.*, Hum.

Gene Ther., 1996, 7:913-919; Rasko *et al.*, Proc. Natl. Acad. Sci. USA, 1999, 96:2129-2134). However, in practice it is highly undesirable to co-culture the target cells with the producer cells as this is not considered a Good Manufacturing Process. Indeed, one cannot regard the results of this experiment with high confidence. Using this approach, the level
 5 of transduction was only the same as with amphotropic viral particles, which indicates that the method does not correlate with previously noted *in vitro* data.

Thus, there is a clear need in the art for more efficient, and thus cost effective, *ex vivo* transduction of stem cells. There is a further need for *ex vivo* transduction of stem cells without inducing stem cell differentiation. The present
 10 invention advantageously addresses these and other needs in the art.

The citation of any reference herein should not be construed as an admission that such reference is available as prior art to the instant invention.

15

SUMMARY OF THE INVENTION

The invention provides a highly efficient method for transducing stem cells with a vector particle, particularly a retroviral vector particle, containing a gene of interest, e.g., in a modified retroviral genome. This method comprises contacting target cells with
 20 vector particles pseudotyped with feline endogenous virus RD114 envelope protein (including binding fragments and fusion proteins) and containing a gene of interest. A feature of the invention is that the vector particles are substantially free of producer cells and producer cell supernatant. Thus, high efficiency transduction of the invention is consistent with Good Manufacturing Practices and avoids conditions that are likely to
 25 induce stem cell differentiation. The advantages of the invention can be achieved by using artificial RD114 pseudotyped particles, by pre-adsorbing the retroviral particles onto a surface that promotes adherence of the retroviral particles, or by freeing the viral particles of producer cells and producer cell supernatant by ultracentrifugation, or both of the latter two.

30

According to the disclosed method, the target stem cells are pre-stimulated by treatment with signaling molecules selected from the group consisting of cytokines, growth factors and phytohemagglutinin.

progenitor cells derived from primitive hematopoietic cells. Such cells can be cord blood cells, mobilized peripheral blood cells, bone marrow cells, and liver cells. Preferably they are CD34⁺ cells; more preferably they are CD34⁺,CD38⁻ cells.

The invention further provides a population of stem cells transduced with
 5 vector particles, particularly retroviral vector particles, pseudotyped with feline endogenous virus RD114 envelope protein and containing a gene of interest. The population of stem cells are substantially undifferentiated. Upon engraftment of the transduced stem cells contacted one time with the retroviral particles into a host, greater than 10% of the transduced cells can express the gene of interest. In a more specific
 10 embodiment, greater than about 40% of the transduced cells express the gene of interest. These are remarkable results.

In addition, the invention provides a method for introducing a gene of interest into a host. This method comprises introducing the population of transduced stem cells of the invention into a host, *e.g.*, a human (and the stem cells are human stem cells), a
 15 non-human host (when the stem cells are autologous or syngenic to the non-human host), or an immunodeficient animal (and the stem cells are human stem cells). Preferably, greater than about 40% of the engrafted transduced stem cells express the gene of interest.

Thus, the invention further provides a non-human animal engrafted with the transduced stem cells of the invention, such as an immunodeficient mouse or a monkey.

20 The invention further provides a method of treating a disease or disorder, which method comprises administering to a patient a therapeutically effective dose of the population of transduced stem cells of the invention, wherein the gene of interest is a therapeutic gene.

A kit comprising retroviral vector particles pseudotyped with feline
 25 endogenous virus RD114 envelope protein and containing a gene of interest their genome pre-adsorbed onto a surface that promotes adherence of the retroviral particles, wherein the retroviral vector particles are substantially free of producer cells and producer cell supernatant, is also provided.

Also, a method for preparing the kit is disclosed. This method comprises
 30 contacting the surface with the retroviral vector particles for a sufficient period of time to permit adherence of the retroviral particles to the surface, and removing supernatant in which the retroviral particles were suspended from the surface. The invention

advantageously permits storing the retroviral particles adsorbed onto the surface, *e.g.*, at -70°C.

DETAILED DESCRIPTION OF THE INVENTION

5 The present invention relates to a method for efficiently introducing exogenous genes into stem cells which, in turn, leads to introducing the exogenous genes into cell lineages derived from these stem cells. Moreover, the high efficiency transduction achieved in accordance with this invention can be achieved without inducing stem cell differentiation. The invention is based, in part, on the discovery that RD114
10 pseudotyped retroviral vectors efficiently transduce hematopoietic stem cells due to unique properties and stability of the vector preparations. These properties include the unexpected ability to highly concentrate RD114 envelope pseudotyped retroviral vectors by adsorbing it on a surface or by ultracentrifugation, thus permitting more effective transduction free of producer cells and producer cell supernatant. Thus, advantages of the
15 present invention are due to the unique, and previously unknown (as well as unexpected) properties of RD114 pseudotyped retroviruses.

 The term “vector particle” refers to a retroviral vector particle or an artificial particle, such as a liposome, protein matrix, or other gene transfer particle (described in greater detail, *infra*) that contains genetic material for transfer into and expression in a cell.
20 The genetic material can be a modified retroviral genome or a recombinant DNA or RNA construct.

 As used herein, “retroviral vector” and “retroviral vector particle” refer to a modified retroviral genome that contains a heterologous or exogenous gene of interest within the retroviral packaging signals, that has been incorporated within a retroviral
25 particle assembled by a producer cell. The retroviral particles of the present invention contain an RD114 envelope (env) protein; thus, they are RD114-pseudotyped retroviral particles.

 As one of ordinary skill in the art would appreciate, the term “pseudotyped with feline endogenous virus RD114 envelope protein” means that the vector particle
30 contains an RD114 protein, or the N-terminal segment of the protein involved in binding, so that the vector particle demonstrates binding properties of RD114. The RD114 protein or binding portion can be included in a fusion construct.

By “highly efficient”, the present invention contemplates significant improvements over the efficiency of stem cell transduction previously reported. It should be noted that efficiency as used herein refers to transduction of stem cells capable of expressing the transgene (the gene of interest) *in vivo*; *in vitro* transduction efficiencies can be very high, for example on the order of 90%. The only level of transduction efficiency that has any relevance is long term marking in hosts, particularly humans and large animals, like the rhesus monkey. However, the highest level of transduction efficiency reported *in vivo* (*i.e.*, after engraftment of the cells transduced *ex vivo*), has been reported to be less than 10% per number of exposures to (*i.e.*, times contacted with) transducing retroviral vectors, as discussed above. The best level of transduction efficiency of which the inventors are aware in a relevant model is no more than 10%. In a specific embodiment, the present invention permits transduction efficiency of about 40% after a single transduction step (contacting the target stem cells with the retroviral vector preparation one time). When considered at the level of the best single animal, the prior art has reported 71% engraftment (in mice), compared to the present invention with an efficiency exceeding 90% in individual animals.

High efficiencies of the invention are achieved by contacting the stem cells with retroviral vector particles pre-adsorbed onto a surface. In a preferred embodiment, the single exposure of stem cells to the pre-adsorbed retroviral particles results in very high levels of transduction efficiency. Higher levels still can be achieved by supplying additional retroviral particles. In order to preserve the phenotype of the target stem cells, the amount of producer cell supernatant added with this second exposure to retroviral particles should be minimized. An advantage of the present invention is the discovery that the RD114 env protein permits concentration of the retroviral particles by ultracentrifugation. Thus, a second exposure or contacting step of the stem cells with retroviral particles is preferably achieved with ultracentrifuge-concentrated retroviral particles.

The term “*ex vivo*” is used herein to refer to transduction of target stem cells with a retroviral vector of the invention in a culture system outside of the host, followed by administration of the transduced stem cells into the host, *i.e.*, engraftment of the transduced stem cells. This term is used in contrast to *in vitro*, which only refers to transduction and maintenance of target cells in tissue culture, and *in vivo*, in which the

vector is administered to the host animal for transduction of the endogenous cells.

As used herein, the term "stem cells" includes but is not limited to hematopoietic stem cells, neural stem cells, mesenchymal (particularly muscular) stem cells, and liver stem cells. Stem cells are capable of repopulating tissue(s) *in vivo*.

- 5 Hematopoietic stem cells are progenitor cells derived from primitive human hematopoietic cells. In a specific embodiment, *infra*, for gene transfer into hematopoietic stem cells, the target cells are selected from the group consisting of cord blood cells, mobilized peripheral blood cells, bone marrow cells, and liver cells. The cells best able to repopulate hematopoietic tissues are CD34⁺ cells, and preferably CD34⁺,CD38⁻ cells. "Mesenchymal stem cells" refer to the cells isolated from connective tissue, including muscle and dermis, which have the ability to differentiate into several phenotypes of the mesodermal lineage, including cartilage and bone (*see* U.S. Patent Nos. 5,906,934, 5,827,735, and 5,486,359). The term "target" is used to indicate that the retroviral vector is intended to transduce the cells.

- 15 A stem cell "population" refers to the stem cells following contact with and transduction by the retroviral vector; because not every stem cell is transduced or transduced effectively (such that it expresses the transgene after engraftment), the cells constitute a heterogeneous population, hence use of that term. While it is possible to isolate or purify the transduced cells (*e.g.*, by including a selection marker in the retroviral vector genome), that is not necessary in the practice of the invention (and, indeed, to the extent that extraneous genes and gene products, including selection genes, are undesirable when the transduced cells are engrafted into a human host for a gene therapy, such markers are preferably avoided).

- 25 The novel gene transfer method of the instant invention involves the use of the retroviral vector (also referred to herein as "retroviral vector particles") pseudotype with feline endogenous virus RD114 envelope protein. As disclosed in specific embodiments, these RD114 pseudotyped retroviral vector particles are generated from producer cells which comprise: (i) a polynucleotide encoding a minimal *gag-pol* open reading frame (ORF) and expressing gag and pol proteins; (ii) a polynucleotide encoding a minimal ORF of feline endogenous virus RD114 envelope protein (*env*); (iii) a retroviral vector including a 5' LTR, a 3' LTR, a packaging signal, and a gene of interest encoding a protein or polypeptide of interest under control of an appropriate expression control
- 30

sequence ("vector genome"). It follows that the retroviral vector particles employed in the present invention contain in their genome only the heterologous coding sequences or sequences and the signals needed for particle assembly (packaging). These vectors can efficiently replicate to produce infective virus only in the producer (helper) cells (which
 5 supply gag, pol and env proteins), and undergo a single cycle of replication and insertion upon infecting a target cell.

The retroviral vector particles are "substantially free of factors that induce stem cell differentiation" when they are substantially free of producer cells and producer cell supernatant, or when the packaging cells do not themselves produce differentiation
 10 factors that affect the stem cells, or when the producer cell culture fluid is treated to neutralize or remove any such factors, *e.g.*, with inhibitory antibodies or by immunoprecipitation. The retroviral vector particles are "substantially free of producer cells and producer cell supernatant" when they have been isolated from the producer cell culture in which they were produced. The term "substantially" with respect to the
 15 producer cells means that the presence of producer cells cannot be detected by microscopy, more preferably by immunoassay, more preferably still by nucleic acid hybridization (Northern or Southern hybridization), and most preferably by nucleic acid amplification, *e.g.*, by polymerase chain reaction (PCR). The term substantially with respect to culture fluid means that the amount of differentiation-inducing factor in producer cell culture fluid
 20 present is too low to cause the stem cells to differentiate, proliferate, die, or undergo any other undesirable outcome. In a specific embodiment, substantially free of producer cell culture fluid means that, when contacted with the target stem cells, less than about 10% of the culture fluid in which the stem cells are suspended is producer cell culture fluid; preferably less than about 1%; and more preferably less than about 0.1%.

25 In specific embodiments, the producer cells of the present invention are derived from HT1080 human sarcoma cell line (ATCC CCL-121), 293T human embryonic kidney cell line (ATCC CRL-1573), or NIH 3T3 mouse fibroblast cell line (ATCC CRL-1658), to mention a few possibilities. The retroviral vector particles can be found in the producer cell culture fluid, which is the culture medium in which the producer
 30 cells grow and effectively permit retroviral vector replication and production.

According to the invention, the retroviral vector particles can be oncoviral particles or lentiviral particles. In a specific embodiment, the RD114 pseudotyped

oncoviral vector genome is derived from murine leukemia virus (MLV). In another specific embodiment, the RD114 pseudotyped oncoviral vector genome is derived from mouse stem cell virus (MSCV). The present invention also discloses the generation of HIV-1-based RD114 pseudotyped lentiviral vector particles. In each of these cases, the
5 key element is using an RD114 *env* gene in the producer cell.

As used herein, the term "isolated" means that the referenced material is free of components found in the natural environment in which the material is normally found. In particular, isolated biological material is free of cellular components. In the case of nucleic acid molecules, an isolated nucleic acid includes a PCR product, an
10 isolated mRNA, a cDNA, or a restriction fragment. In another embodiment, an isolated nucleic acid is preferably excised from the chromosome in which it may be found, and more preferably is no longer joined to non-regulatory, non-coding regions, or to other genes, located upstream or downstream of the gene contained by the isolated nucleic acid molecule when found in the chromosome. In yet another embodiment, the isolated nucleic
15 acid lacks one or more introns. Isolated nucleic acid molecules can be inserted into plasmids, cosmids, artificial chromosomes, and the like. Thus, in a specific embodiment, a recombinant nucleic acid is an isolated nucleic acid. An isolated protein may be associated with other proteins or nucleic acids, or both, with which it associates in the cell, or with cellular membranes if it is a membrane-associated protein. An isolated organelle,
20 cell, or tissue is removed from the anatomical site in which it is found in an organism. An isolated material may be, but need not be, purified.

The term "purified" as used herein refers to material that has been isolated under conditions that reduce or eliminate unrelated materials, *i.e.*, contaminants. For example, a purified protein is preferably substantially free of other proteins or nucleic
25 acids with which it is associated in a cell; a purified nucleic acid molecule is preferably substantially free of proteins or other unrelated nucleic acid molecules with which it can be found within a cell. As used herein, the term "substantially free" is used operationally, in the context of analytical testing of the material. Preferably, purified material substantially free of contaminants is at least 50% pure; more preferably, at least 90% pure, and more
30 preferably still at least 99% pure. Purity can be evaluated by chromatography, gel electrophoresis, immunoassay, composition analysis, biological assay, and other methods known in the art.

The phrase "pharmaceutically acceptable", whether used in connection with the pharmaceutical compositions of the invention or vaccine compositions of the invention, refers to molecular entities and compositions that are physiologically tolerable and do not typically produce untoward reaction, such as gastric upset, dizziness and the like, when administered to a human. Preferably, as used herein, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the compound is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water or aqueous solution saline solutions and aqueous dextrose and glycerol solutions are preferably employed as carriers, particularly for injectable solutions. Suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin, 18th Edition.

The term "about" or "approximately" will be known to those skilled in the art in light of this disclosure. Generally it means that a particular value can range within an acceptable error for the type of measurement from which the value is obtained. For example, the term can mean within 20%, more preferably within 10%, and more preferably still within 5% of a given value. Alternatively, especially in biological systems, the term "about" preferably means within about a log (*i.e.*, an order of magnitude), preferably within a factor of five and more preferably within a factor of two, of a given value, depending on how quantitative the measurement.

25

Molecular Biology - Definitions

In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, *e.g.*, Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual*, Second Edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (herein "Sambrook *et al.*, 1989"); *DNA Cloning: A Practical Approach*, Volumes I and II (D.N. Glover ed. 1985); *Oligonucleotide Synthesis* (M.J. Gait ed. 1984); *Nucleic Acid*

(D.N. Glover ed. 1985); *Oligonucleotide Synthesis* (M.J. Gait ed. 1984); *Nucleic Acid Hybridization*, B.D. Hames & S.J. Higgins eds. (1985); *Transcription And Translation*, B.D. Hames & S.J. Higgins, eds. (1984); *Animal Cell Culture*, R.I. Freshney, ed. (1986); *Immobilized Cells And Enzymes*, IRL Press, (1986); B. Perbal, *A Practical Guide To*
 5 *Molecular Cloning* (1984); F.M. Ausubel *et al.* (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc. (1994).

A "coding sequence" or a sequence "encoding" an expression product, such as a RNA, polypeptide, protein, or enzyme, is a nucleotide sequence that, when expressed, results in the production of that RNA, polypeptide, protein, or enzyme, *i.e.*, the nucleotide
 10 sequence encodes an amino acid sequence for that polypeptide, protein or enzyme. A coding sequence for a protein may include a start codon (usually ATG) and a stop codon.

The term "gene", also called a "structural gene" means a DNA sequence that codes for or corresponds to a particular sequence of amino acids which comprise all or part of one or more proteins, and may or may not include regulatory DNA sequences, such
 15 as promoter sequences; that determine for example the conditions under which the gene is expressed. The transcribed region of a gene can include 5'- and 3'-untranslated regions (UTRs) and introns in addition to the translated (coding) region. As used herein, the term "gene" in conjunction with a vector means a coding sequence operatively associated with an expression control sequence; it can therefore be an artificial construct.

A "promoter sequence" is a DNA regulatory region capable of binding
 20 RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate
 25 transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site (conveniently defined for example, by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase.

Promoters that may be used to control gene expression include, but are not
 30 limited to, cytomegalovirus (CMV) promoter (U.S. Patent Nos. 5,385,839 and 5,168,062), the SV40 early promoter region (Benoist and Chambon, *Nature*, 1981, 290:304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, *et*

Natl. Acad. Sci. U.S.A., 1981, 78:1441-1445), the regulatory sequences of the metallothionein gene (Brinster *et al.*, Nature, 1982, 296:39-42); prokaryotic expression vectors such as the β -lactamase promoter (Villa-Komaroff, *et al.*, Proc. Natl. Acad. Sci. U.S.A., 1978, 75:3727-3731), or the *tac* promoter (DeBoer, *et al.*, Proc. Natl. Acad. Sci. U.S.A., 1983, 80:21-25); *see also* "Useful proteins from recombinant bacteria" in Scientific American, 1980, 242:74-94; promoter elements from yeast or other fungi such as the Gal 4 promoter, the ADC (alcohol dehydrogenase) promoter, PGK (phosphoglycerol kinase) promoter, alkaline phosphatase promoter; and transcriptional control regions that exhibit hematopoietic tissue specificity, in particular: beta-globin gene control region which is active in myeloid cells (Mogam *et al.*, Nature, 1985, 315:338-340; Kollias *et al.*, 1986, Cell 46:89-94), hematopoietic stem cell differentiation factor promoters, erythropoietin receptor promoter (Maouche *et al.*, Blood, 1991, 15:2557), etc.

A coding sequence is "under the control" of or "operably associated with" transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then trans-RNA spliced (if it contains introns) and translated into the protein encoded by the coding sequence.

The terms "express" and "expression" mean allowing or causing the information in a gene or DNA sequence to become manifest, for example producing a protein by activating the cellular functions involved in transcription and translation of a corresponding gene or DNA sequence. A DNA sequence is expressed in or by a cell to form an "expression product" such as an mRNA or a protein. The expression product itself, *e.g.* the resulting mRNA or protein, may also be said to be "expressed" by the cell. An expression product can be characterized as intracellular, extracellular or secreted. The term "intracellular" means something that is inside a cell. The term "extracellular" means something that is outside a cell. A substance is "secreted" by a cell if it appears in significant measure outside the cell, from somewhere on or inside the cell. "Conditions that permit expression", *in vitro* are culture conditions of temperature (generally about 37°C), humidity (humid atmosphere), carbon dioxide concentration to maintain pH (generally about 5% CO₂ to about 15% CO₂), pH (generally about 7.0 to 8.0, preferably 7.5), and culture fluid components, that depend on host cell type. *In vivo*, the conditions that permit expression are primarily the health of the animal, which depends on adequate nutrition, water, habitation, and environmental conditions (light-dark cycle, temperature,

humidity, noise level). In either system, expression may depend on a repressor or inducer control system, as well known in the art.

The term "transfection" means the introduction of a "foreign" (*i.e.* extrinsic or extracellular) gene, DNA or RNA sequence into a host cell, so that the host cell will
 5 express the introduced gene or sequence to produce a desired substance, typically a protein or enzyme coded by the introduced gene or sequence. The introduced gene or sequence may also be called a "cloned" or "foreign" gene or sequence, may include regulatory or control sequences, such as start, stop, promoter, signal, secretion, or other sequences used by a cell's genetic machinery. The gene or sequence may include nonfunctional sequences
 10 or sequences with no known function. A host cell that receives and expresses introduced DNA or RNA has been "transfected" and is a "transfectant" or a "clone." The DNA or RNA introduced to a host cell can come from any source, including cells of the same genus or species as the host cell, or cells of a different genus or species.

The terms "vector", "cloning vector" and "expression vector" mean the
 15 vehicle by which a DNA or RNA sequence (*e.g.* a foreign gene) can be introduced into a host cell, so as to transform the host and promote expression (*e.g.* transcription and translation) of the introduced sequence. Vectors include plasmids, phages, viruses, etc.; they are discussed in greater detail below.

Retroviral vectors typically comprise the RNA of a transmissible agent, into
 20 which a heterologous sequence encoding a protein of interest is inserted. Typically, the retroviral RNA genome is expressed from a DNA construct. A common way to insert one segment of DNA into another segment of DNA involves the use of enzymes called restriction enzymes that cleave DNA at specific sites (specific groups of nucleotides) called restriction sites. A "cassette" refers to a DNA segment that can be inserted into a
 25 vector or into another piece of DNA at a defined restriction site. Preferably, a cassette is an "expression cassette" in which the DNA is a coding sequence or segment of DNA that codes for an expression product that can be inserted into a vector at defined restriction sites. The cassette restriction sites generally are designed to ensure insertion of the cassette in the proper reading frame. Generally, foreign DNA is inserted at one or more
 30 restriction sites of the vector DNA, and then is carried by the vector into a host cell along with the transmissible vector DNA. A segment or sequence of DNA having inserted or added DNA, such as an expression vector, can also be called a "DNA construct." A

common type of DNA construct is a "plasmid" that generally is a self-contained molecule of double-stranded DNA, usually of bacterial origin, that can readily accept additional (foreign) DNA and which can be readily introduced into a suitable producer cell. A plasmid vector often contains coding DNA and promoter DNA and has one or more

5 restriction sites suitable for inserting foreign DNA. A large number of vectors, including plasmid and fungal vectors, have been described for replication and/or expression in a variety of eukaryotic and prokaryotic hosts. Non-limiting examples include pKK plasmids (Amersham Pharmacia Biotech), pUC plasmids, pET plasmids (Novagen, Inc., Madison, WI), pRSET or pREP plasmids (Invitrogen, San Diego, CA), or pMAL plasmids (New

10 England Biolabs, Beverly, MA), and many appropriate host cells, using methods disclosed or cited herein or otherwise known to those skilled in the relevant art. Recombinant cloning vectors will often include one or more replication systems for cloning or expression, one or more markers for selection in the host, *e.g.* antibiotic resistance, and one or more expression cassettes. A "retroviral plasmid vector" means a plasmid which

15 includes all or part of a retroviral genome including 5' and 3' retroviral long-term repeat (LTR) sequences, a packaging signal (.psi.), and may include one or more polynucleotides encoding a protein(s) or polypeptide(s) of interest, such as a therapeutic agent or a selectable marker. Such retroviral plasmid vectors are described, *e.g.*, in U.S. Patent No. 5,952,225 (column 4, line 5 to column 5, line8), which is specifically incorporated herein

20 by reference.

Synthetic vector particles can be prepared using lipofection technology, optionally with other transfection facilitating agents (peptides, polymers, etc.). Synthetic cationic lipids can be used to prepare liposomes for *in vivo* transfection of a gene (Felgner, et. al., Proc. Natl. Acad. Sci. USA 1987, 84:7413-7417; Felgner and Ringold, Science

25 1989, 337:387-388; see Mackey, *et al.*, Proc. Natl. Acad. Sci. USA 1988, 85:8027-8031; Ulmer *et al.*, Science 1993, 259:1745-1748). Useful lipid compounds and compositions for transfer of nucleic acids are described in PCT Publications WO95/18863 and WO96/17823, and in U.S. Patent No. 5,459,127. Lipids may be chemically coupled to

RD114 molecules for the purpose of targeting (see Mackey, et. al., *supra*), or by insertion

30 of an RD114 polypeptide construct into the lipid bilayer, *i.e.*, by analogy to a transmembrane protein.

The term "host" or "host animal" means any animal that is selected,

modified, engrafted, or manipulated in any way, for the production of the protein of interest (expressed by the gene of interest) in the host. Non-human animal hosts can further be used for screening or other assays, as described *infra*. Human hosts can be used to study the distribution and fate of engrafted stem cells, *e.g.*, carrying a marker gene.

5 The term "heterologous" refers to a combination of elements not naturally occurring. For example, heterologous DNA refers to DNA not naturally located in the cell, or in a chromosomal site of the cell. A heterologous expression regulatory element is such an element operatively associated with a different gene than the one it is operatively associated with in nature. In the context of the present invention, a coding sequence of
10 interest is heterologous to the retroviral vector RNA in which it is inserted for expression, and it is heterologous to a host cell or animal containing such a vector in which it is expressed, *e.g.*, a CHO cell, a mouse, a monkey, or a human.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler *et al.*, Cell, 1977, 11:223),
15 hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, Proc. Natl. Acad. Sci. USA, 1962, 48:2026), and adenine phosphoribosyltransferase (Lowy *et al.*, Cell, 1980, 22:817) genes can be employed in tk-, hgp^rt-, or ap^rt- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler *et al.*, Proc. Natl. Acad. Sci. USA,
20 1980, 77:3567; O'Hare *et al.*, Proc. Natl. Acad. Sci. USA, 1981, 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, Proc. Natl. Acad. Sci. USA, 1981, 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin *et al.*, J. Mol. Biol., 1981, 150:1); and hyg^r, which confers resistance to hygromycin (Santerre *et al.*, Gene, 1984, 30:147).

25 One aspect of the invention involves the use of a retroviral vector genome expressing a bicistronic transcript encoding the gene of interest and a selectable marker gene, wherein the selectable marker gene is arranged downstream of the stop codon of the gene of interest and is separated from it by an internal ribosome entry site to insure that it is expressed from the corresponding mRNA as a result of translation reinitiation.
30 According to the invention, the inclusion of a selectable marker gene allows the efficient selection of producer cells and transduced target cells. Thus, in one embodiment, the present invention discloses the use of a selectable marker gene encoding a drug-resistant

variant of human dehydrofolate reductase which confers the resistance to trimetrexate. In another embodiment, the present invention discloses the use of retroviral vectors which do not encode a selectable marker, but only the gene(s) of interest.

5

Use of the Vector Particles

According to the invention, the gene of interest carried by the RD114 can be any gene pseudotyped vector particle. In a preferred embodiment, the gene of interest is a therapeutically relevant gene. The non-limiting examples of such genes include genes encoding wild-type proteins missing in mutant cells (*e.g.*, factors VII and IX, tumor suppressor genes, etc.) and genes involved in drug resistance or anti-viral resistance (*e.g.*, MDR, ribozymes, antisense RNAs, anti-viral proteases, etc.). Examples of therapeutic genes include polynucleotides encoding tumor necrosis factor (TNF) genes, such as TNF- α ; genes encoding interferons such as Interferon- α , Interferon- β , and Interferon- γ ; genes encoding interleukins such as IL-1, IL- β , and Interleukins 2 through 14; genes encoding GM-CSF; genes encoding adenosine deaminase, or ADA; genes which encode cellular growth factors, such as lymphokines, which are growth factors for lymphocytes; genes encoding epidermal growth factor (EGF), and keratinocyte growth factor (KGF); genes encoding soluble CD4; Factor VIII; Factor IX; cytochrome b; glucocerebrosidase; T-cell receptors; the LDL receptor, ApoE, ApoC, ApoAI and other genes involved in cholesterol transport and metabolism; the alpha-1 antitrypsin (α 1AT) gene; the insulin gene; the hypoxanthine phosphoribosyl transferase gene; the CFTR gene; negative selective markers or "suicide" genes, such as viral thymidine kinase genes, *e.g.*, the Herpes Simplex Virus thymidine kinase gene, the cytomegalovirus virus thymidine kinase gene, and the varicella-zoster virus thymidine kinase gene; Fc receptors for antigen-binding domains of antibodies, antisense sequences which inhibit viral replication, such as antisense sequences which inhibit replication of hepatitis B or hepatitis non-A non-B virus; antisense c-myc oligonucleotides; anti-tumor protein intracellular antibodies; and antioxidants such as, but not limited to, manganese superoxide dismutase (Mn-SOD), catalase, copper-zinc-superoxide dismutase (CuZn-SOD), extracellular superoxide dismutase (EC-SOD), and glutathione reductase; tissue plasminogen activator (tPA); urinary plasminogen activator (urokinase); hirudin; the phenylalanine hydroxylase gene; nitric oxide synthetase; vasoactive peptides; angiogenic peptides; the dopamine gene; the

nitric oxide synthetase; vasoactive peptides; angiogenic peptides; the dopamine gene; the dystrophin gene; the β -globin gene; the α -globin gene; the HbA gene; protooncogenes such as the ras, src, and bcl genes; tumor-suppressor genes such as p53 and Rb; the LDL receptor; the heregulin- α protein gene, for treating breast, ovarian, gastric and endometrial cancers; monoclonal antibodies specific to epitopes contained within the β -chain of a T-cell antigen receptor; the multidrug resistance (MDR) gene; polynucleotides encoding ribozymes; antisense polynucleotides; genes encoding secretory peptides which act as competitive inhibitors of angiotensin converting enzyme, of vascular smooth muscle calcium channels, or of adrenergic receptors, and polynucleotides encoding enzymes which break down amyloid plaques within the central nervous system. It is to be understood, however, that the scope of the present invention is not to be limited to any particular therapeutic agent. In one of the embodiments, the invention also discloses the use of a gene of interest encoding a marker gene, such as enhanced green fluorescent protein (EGFP). Such marker genes are particularly useful for tracking the fate of transduced cells, *e.g.*, during clinical development of a gene therapy based on the vectors and methods of the invention.

The present invention also discloses a method for somatic gene therapy which involves introducing a gene of interest contained within the vector particle, such as in a retroviral genome, into human repopulating stem cells followed by introducing these cells into a human host. Such therapies can be tested on the animal models described herein. In accord with the present invention, an animal model system is described for use in retroviral gene therapy studies, which employs an efficient and rapid protocol for the introduction of exogenous genes.

Because the vector particles of the invention are highly efficient at transducing stem cells, they are particularly attractive for somatic gene therapy. Gene therapy can be used to treat any disorders characterized by a defect in a single gene, or which can be treated by producing expression of a gene, even though the cause of the disease is a multi-genic defect, or some non-genetic cause, such as an infection or autoimmunity. In particular, diseases that can be treated with the invention, in addition to those set forth in the background of the invention or that are obvious from the foregoing list of therapeutic genes, are hematopoietic disease, neural disease, joint-related disease, muscular disease, and liver disease. In particular, joint-related diseases include

cells may be beneficial.

The present invention yet further discloses a method for monitoring the efficiency of the stem cell-mediated gene transfer based on detecting the presence of the genes (or the expression products) of the vector in various stem cell-derived lineages of the host. As disclosed in a specific embodiment, the presence of the genes of the vector can be monitored by any of the methods known in the art, but most preferably by PCR (using retroviral-specific primers) and/or by detecting the polypeptide product of the vector-encoded gene(s) (*e.g.*, using immunochemical or fluorescence assays).

The disclosed novel highly efficient method of vector particle-mediated transduction of stem cells preferably comprises the steps of : (i) optionally inducing the proliferation of a target stem cell by pre-stimulation; and (ii) incubating the pre-stimulated target stem cell with a vector particle containing a gene of interest and pseudotyped with feline endogenous virus RD114 envelope protein. While traditionally, the transduced cells are expanded in culture, the present invention advantageously omits this step, and thus avoids inducing stem cell differentiation. In a less preferred embodiment, the target stem cell is pre-stimulated with cytokines and/or growth factors. These cytokines and growth factors can be selected from the group consisting of but not limited to: stem cell factor, Flt-3 ligand, interleukin-2, interleukin-3, interleukin-6, and phytohemagglutinin. According to the method of the instant invention, the optimal efficiency of retroviral-stimulated stem cell transduction requires using target cell density around $1-5 \times 10^4$ cells/cm² and is attained upon pre-stimulating target cells for 24-48 hours.

As pointed out above, the method of the present invention benefits from the use of RD114 pseudotyped retroviral vector particles substantially free of producer cells and producer cell supernatant. Such retroviral particles can be either pre-adsorbed on retronectin coated plates or concentrated by ultracentrifugation. As disclosed in a specific embodiment, HT1080-derived producer cells (and potentially other kinds of producer cells) contain in their medium an unidentified substance that induces phenotypic change in the target cells, leading to their elimination upon engraftment in the immunodeficient host. According to the method of the instant invention, retronectin-pre-adsorbed and ultrafiltration-concentrated retroviral particles and synthetic vector particles that contain RD114 binding polypeptides are free of contaminating and potentially harmful substances present in the culture media of producer cells, leading to the efficient transduction and

engraftment of the target cells without any change in their phenotype. In a preferred embodiment, the disclosed method involves producing retronectin plates with pre-adsorbed retroviral vector particles as a part of a transduction kit which can be stored at low temperatures for long periods of time.

5 As exemplified in examples below, cells can be transduced by the vector particles of the invention by direct contact in tissue culture. For example, in a specific embodiment, cells are exposed to pre-loaded vector (adsorbed to surfaces) at 48 hours in culture. Alternatively, concentrated vector can be added to these cells. In yet another alternative, the cells at 48 hours in culture are contacted with pre-loaded vector, and
10 concentrated vector is added at 72 hours. While this latter addition results in contact or exposure to producer cell supernatant, the total amount of supernatant is limited. This is because ultracentrifugation concentrates the virus 50 to 100-fold, although the concentrated virus is in approximately 50% solution of supernatant. Adding this to the media containing the target stem cells at a 10% volume means exposing the cells to 5%
15 producer cell supernatant. Presumably, as a result of centrifugation, most of the adverse proteins do not concentrate due to their small size. Thus, the target stem cells are exposed to less than 10% of the concentrated producer cell fluid, which is likely to contain less than a proportional amount of unwanted proteins.

20

Pre-adsorption of Retroviral Vector Particles

As noted above, the retroviral particles can be pre-adsorbed onto a surface that promotes adherence of the retroviral particles. Optimal transduction using RD114-pseudotyped vectors of stem cells can be obtained by preloading/adsorption of vectors on
25 surfaces, *e.g.*, retronectin-coated culture dishes.

Suitable surfaces include, but are by no means limited to, plastic, glass, polymer particles (such as SEPHADEX and SEPHAROSE), gels, and the like. Preferred plastic surfaces are tissue culture vessels, flasks, or plates.

Preferably, the surface is coated with an adherence promoting agent. Such
30 adherence promoting agents are typically biomolecules to which an RD114 pseudotyped virus adheres. Adherence can be detected by demonstrating the presence of more RD114 pseudotyped retroviral particles adhering to a surface in the presence relative to the

absence of an adherence promoting agent. In specific embodiments, the adherence promoting agent is retronectin, fibronectin, or polylysine; retronectin is exemplified *infra*.

To adsorb the retroviral particles, retroviral particle-containing supernatant from a producer cell culture is contacted with the solid surface for a time and under
 5 conditions of temperature, pH, humidity, etc. that permit adsorption to occur. Once the particles are adsorbed, the supernatant can be decanted or otherwise removed. If desired, the adsorbed particles can be washed to remove additional impurities, *e.g.*, proteins from the supernatant, although the washing step should be done under mild conditions so as not to remove the pre-adsorbed retroviral particles.

10

Pre-adsorbed Transduction Kits

Having shown that pre-adsorption of RD114 pseudotyped retroviral particles greatly increases transduction efficiency, it was further discovered that the pre-adsorbed particles could be stored for later use, thus permitting development of kits. Thus,
 15 the invention encompasses a service for producing high efficiency retroviral vector particles.

A kit of the invention comprises retroviral vector particles pseudotyped with feline endogenous virus RD114 envelope protein and containing a gene of interest their genome pre-adsorbed onto a surface that promotes adherence of the retroviral
 20 particles. As noted above, the retroviral vector particles are substantially free of producer cells and producer cell supernatant. Preferred kits employ plastic tissue culture containers (plates or vessels) as the surface to which the retroviral particles are adsorbed. The kits can be stored prior to use, preferably under conditions that preserve the transduction capability of the retrovirus particles, *e.g.*, about 50%, preferably greater than about 75%,
 25 and more preferably greater than about 90% of the transduction potential of the retroviral particles at the time the kit is prepared. In a specific embodiment, the adsorbed particles are stored at -70°C. Experiments showed that the transduction efficiency was maintained after 48 hours of storage, one week of storage, and for longer periods under these conditions.

30

Accordingly, the invention provides a method for preparing such a kit. This method comprises contacting the surface with the retroviral vector particles for a sufficient period of time to permit adherence of the retroviral particles to the surface, and

removing supernatant in which the retroviral particles were suspended from the surface. The surface can be washed.

Thus, a vector production service can be implemented, in which a gene of interest is provided by a customer to the service lab, who introduce it into an RD114-pseudotyped retroviral vector. The recombinant vectors are then adsorbed onto a suitable surface, *e.g.*, a retronectin-coated tissue culture dish, and provided to the customer. The customer can use this kit, containing retroviral vectors that express the customer's gene of interest, to transduce stem cells.

Transduced Animal Models

Because of the high efficiency of transduction achieved by the present invention, it is possible to create animal models for human conditions by engrafting non-human animals with autologous or human stem cells transduced in accordance with the invention. Thus, the invention provides such non-human animals. Preferably, if the stem cells are human stem cells the animal is immunodeficient, so that it does not reject the engrafted cells. However, in pre-clinical studies, engraftment with autologous cells is desired. The present inventors have successfully engrafted monkeys with the same high level of efficiency found with immunodeficient mice.

The animal models can be used to study the fate of marker-gene containing transduced stem cells. Animal models in which the animals are engrafted with human stem cells can be used to study the effect of various pharmacological agents on the human cells, or to evaluate the effect of transgene production by the retroviral vectors on the animal physiology.

Suitable animals for these models include, but are by no means limited to, mice, rats, rabbits, hamsters, guinea pigs, and other rodents; dogs, cats, and other domesticated carnivores; sheep, goats, pigs, and other farm animals; and monkeys, chimpanzees, apes, and other primates.

In specific embodiments, the invention discloses a NOD/SCID murine model and an immunodeficient simian model. In one specific embodiment, the present invention discloses the use of RD114 pseudotyped oncoretroviral vector particles for transducing primitive human hematopoietic cells which are then injected into the bone marrow of NOD/SCID mice. According to this embodiment, the injected transduced stem

cells repopulated host bone marrow with around 90% efficiency leading to the introduction of a transgene into various hematopoietic cell lineages. In yet another embodiment, RD114 pseudotyped retroviral particles are used to transduce monkey hematopoietic stem cells, which were then injected into a recipient monkey. According to this embodiment, the attained efficiency of engraftment is around 70%.

EXAMPLES

The invention may be better understood by reference to the following Examples, which is provided by way of exemplification and not limitation.

EXAMPLE 1: Gene Transfer into Primitive Human Hematopoietic Cells Using Retroviral Vector Particles Pseudotyped with the Feline Endogenous Retrovirus (RD114) Envelope Protein

Materials and Methods

Cell lines and primary cell populations. Human hematopoietic cell lines (K562-ATCC CCL 243 and HEL-ATCC TIB 180), mouse fibroblasts (NIH3T3-ATCC CRL 1658), and human embryonic kidney cells (293T-ATCC CRL 1573) were grown in Dulbecco's Modified Eagles Medium (DMEM) supplemented with 10% Fetal Calf Serum (FCS). Human peripheral blood mononuclear cells (PBMC) were recovered from the blood of normal donors by centrifugation on Histopaque-1077 (Sigma). Human umbilical cord blood specimens were obtained from delivered placentas following uncomplicated births at a local delivery center. Mononuclear cell preparations were recovered by centrifugation on Histopaque-1077. The CD34⁺ cells in the cord blood mononuclear cell specimens were purified using a CD34-specific magnetic cell selection system according to instructions provided by the manufacturer (Miltenyi Biotec, Bergisch Gladbach, Germany). CD34⁺, CD38⁻ cells were recovered from the CD34⁺ enriched populations by labeling with anti-CD34 (clone MY10) and anti-CD38 (clone HB-7) monoclonal antibodies (Becton Dickinson, San Jose, CA) conjugated to fluorescein isothiocyanate (FITC) or phycoerythrin (PE), respectively, and sorted for the CD34⁺, CD38⁻ population using a vantage flow cytometer (Becton Dickinson).

Retroviral vector preparations. Vector particles pseudotyped with the feline endogenous virus (RD114) envelope protein were derived by generating producer

5 cells from a packaging cell line designated FLYRD18 (Porter *et al.*, Hum. Gene Ther.,
 1996, 7:913-919) Cossett, J. Virol., 1995, 69:7430; ECACC Accession No. 95091902.
 This packaging cell line had been derived from human sarcoma cells (HT1080) by
 introducing the *gag-pol* genes from murine leukemia virus (MuLV) and the *env* gene
 encoding the envelope protein of the RD114 virus. We derived producer cells from the
 10 FLYRD18 packaging cell line by introducing a vector genome (MGirL22Y) that encodes
 the enhanced green fluorescent protein (EGFP) and a drug resistant variant of human
 dihydrofolate reductase (L22Y) (Allay *et al.*, Nat. Med., 1998, 4:1136-1143; Persons *et al.*,
 Nat. Med., 1998, 4:1201-1205). The reading frames for these proteins are separated by an
 internal ribosomal entry site and transcribed into a bicistronic transcript under the control
 of the mouse stem cell virus (MSCV) long terminal repeat (LTR) (Cheng *et al.*, Gene
 Therapy, 1997, 4:1013-1022). The transduced FLYRD114 cells were selected in
 trimetrexate or sorted for high EGFP expression as previously described (Allay *et al.*,
supra; Persons *et al.*, *supra*). Individual clones were recovered by limiting dilution and
 their capacity for vector production was determined by assaying serial dilutions of
 20 conditioned medium on human HeLa cells and a high titer clone designated
 RD114/MGirL22Y was identified. This methodology was also used to obtain
 amphotropic producer cell populations, AM13/MGirL22Y and AM/MGirL22Y, generated
 from FLYA13 packaging cells (a derivative of HT1080; Cosset *et al.*, 1995, J. Virol.,
 69:7430-7436) and PA317 packaging cells (Miller and Buttimore, 1986, Mol. Cell. Biol.,
 6:2895-2902) (ATTC Accession No. CRL 9078), respectively. GALV pseudotyped
 25 particles were generated by a producer clone derived from PG13 cells (Miller *et al.*, 1991,
 J. Virol., 65:2220-2224) (ATCC Accession No. CRL 10686), and VSV-G pseudotyped
 particles were generated by a producer clone derived from 293T cells (Ory *et al.*, 1996,
 Proc. Natl. Acad. Sci. USA, 93:11400-11406) using the techniques described above.
 30 Vector preparations were screened for replication-competent virus by a marker rescue
 assay using HeLa (ATCC Accession No. CCL 2) or K562 (ATCC Accession No. ATCC
 CCL 243) cells which contained an integrated vector genome encoding neomycin
 resistance (G1NA).

35 Vector particles pseudotyped with the RD114, VSV-G, or amphotropic
 envelope proteins were also generated in 293T cells which had been transfected with
 helper and vector plasmids (Persons *et al.*, Blood Cells Mol. Dis., 1998, 24:167-182). For

RD114 pseudotyped particles, the 293T cells were transfected with a plasmid containing the vector genome (pMGirL22Y), a second plasmid encoding the gag and pol proteins of MuLV (pEQPAM3-E), and a third plasmid encoding the env protein of RD114 feline endogenous retrovirus (pRDF). Between 48-72 hours following transfection, conditioned media was harvested, titered on HeLa cells and screened for replication-competent retrovirus by the marker rescue assay as described above.

In vitro analysis of gene transfer efficiency. All transductions were performed in medium which contained 6 mg/ml of polybrene. Various human and murine cell lines were transduced at specified multiplicities of infection (MOI) based on titers determined on HeLa cells. After an overnight incubation, the vector containing culture medium was removed and fresh medium was added. The cells were harvested 72 hours after transduction and analyzed for EGFP expression by flow cytometry (FACS caliber, Becton Dickinson, San Jose, CA).

Peripheral blood mononuclear cells (PBMC) (lymphocytes) were activated with phytohemagglutinin (PHA, 34mg/ml) and interleukin-2 (IL-2, R&D Systems) at 100U/ml for 48 hours in RPMI-1640 medium supplemented with 10% FCS for 48 hours prior to transduction. These activated human PBMC (2×10^4 /well) were then transduced in 48-well plates that had been coated with retronectin (CH-296, Takara Shuzo, Otsu, Japan) at a concentration of 20mg/cm². Fresh medium containing IL-2 (100U/ml) was used to replace the vector containing medium after overnight incubation and the cells were allowed to expand for 72 hours post-transduction before analysis by flow cytometry. Analysis showed that more than 95% of the cells reacted with an anti-CD3 specific monoclonal antibody (data not shown).

CD34⁺ or CD34⁺, CD38⁻ purified primitive hematopoietic cell populations were cultured in Iscove's Modified Dulbecco's Medium (IMDM) plus 1% bovine serum albumin, human insulin (5mg/ml), human transferrin (100mg/ml), low density lipoproteins (10mg/ml), 0.1mM β -mercaptoethanol, stem cell factor (SCF, 300ng/ml), Flt-3 ligand (300ng/ml), interleukin-3 (IL-3, 10ng/ml) and interleukin-6 (IL-6, 10ng/ml). The cytokines were obtained from R&D Systems, Minneapolis, Minnesota. The cells were incubated for 24 hours at 37°C in 5% CO₂ prior to transduction.

Transduction of the CD34⁺ and CD34⁺, CD38⁻ cells was performed in retronectin coated 48-well plates ($1-2 \times 10^4$ cells per well at a concentration of $1-2 \times 10^5$

cells/ml). When indicated, the retronectin coated wells were pre-loaded with retroviral vector particles by placing 0.5ml/cm² of medium conditioned by producer cells in each well and incubating for 20-30 minutes at room temperature. This medium was then aspirated and a serum-free culture medium (specified above) containing CD34⁺ or CD34⁺, CD38⁻ purified cells was added. When transductions were performed without preloading, serum-containing medium (10% FCS) conditioned by producer cells was added to achieve the specified MOI in amounts up to 40% of the initial culture volume.

Gene transfer efficiency was analyzed by evaluating EGFP expression in transduced cells. After incubation, the CD34⁺ or CD34⁺, CD38⁻ cells were washed in PBS containing 2% heat-inactivated FCS and then stained with mouse anti-CD38 (clone HB-7) and anti-CD34 (clone My10) monoclonal antibodies conjugated to PE or allophycocyanin (APC, Becton Dickinson, San Jose, CA), respectively, washed, and resuspended in PBS containing 2% FCS. Four color flow cytometry was performed and the data were analyzed using the Cell Quest Software Package (Becton Dickinson).

To assay gene transfer into clonogenic progenitors, transduced and control hematopoietic cells were replated after 96 hours of culture into Methocult GF (H4434, Stem Cell Technologies, Vancouver, B.C., Canada) which had been pretreated with 1.2U/ml thymidine phosphorylase at 37°C for two hours. Cultures were established with or without 100nM trimetrexate. At this concentration of trimetrexate, no colonies were formed in the samples containing control (untransduced) cells. Hematopoietic cells were cultured in 35mm plates (1ml of medium per plate) at 37°C in a 5% CO₂ humidified atmosphere for 10-15 days after which the colonies were enumerated.

Analysis of gene transfer into cells that establish human hematopoiesis in immunodeficient (NOD/SCID) mice. Purified CD34⁺ cells were prestimulated in medium containing SCF, Flt-3 ligand, IL-3 and IL-6 as described above for 24 hours at a concentration of 1-2x10⁵ cells/ml. The cells in this medium were then transferred to retronectin coated culture plates to which vector particles had been absorbed (pre-loaded) or to retronectin coated plates without virus. In the later case, vector particles were added in the form of conditioned medium (constituting up to 20% of the culture volume). All cultures were diluted 2-fold with serum-free medium containing cytokines at 48 hours and harvested for injection at 96 hours.

5 The NOD/SCID mice (Jackson Laboratories, Bar Harbor, Maine) were housed in sterile microisolator cages and supplied with sterile food, acidified water and bedding. These six- to eight-week-old mice were used after sublethal irradiation (3-5cGy-¹²⁷Cs source). Each mouse was injected with $1-1.5 \times 10^5$ freshly isolated CD34⁺ cells (>95% purity) or after expansion of this input volume for up to 96 hours in culture. The mice were sacrificed 8-10 weeks following injection and bone marrow cells were harvested for flow cytometric analysis and *in vitro* culturing.

10 Bone marrow cells from animals injected with human cells was subjected to flow cytometric analysis using conjugated antibodies against human surface antigens as follows: 1) CD45-APC to screen human hematopoietic cells; 2) CD19-PE to screen B-lymphocytes; and 3) CD33-PE to screen myeloid cells. These antibodies were obtained from Pharmingen (San Diego, California). $5-10 \times 10^5$ bone marrow cells were mixed with either rat anti-mouse CD16/CD32 Fc block (clone 2.4G2, Pharmingen) or 10% heat-inactivated pooled mouse serum (to block non-specific antibody binding), and then incubated with saturating amounts of one of the conjugated antibodies. Cells from each animal were also stained with appropriate conjugated, isotype matched, control antibodies obtained from Becton Dickinson or Pharmingen. After incubation, cells were resuspended in red cell lysis buffer and washed twice in PBS containing 2% FCS. In all experiments cells stained with the isotype control antibody were used to set the quadrant markers such that the negative quadrant contained at least 97% of the control cells. The percentage of engrafted human cells was determined by CD45 positivity, followed by the determination of lineage marker and EGFP expression in the CD45⁺-gated population.

25 The number of total and trimetrexate-resistant human clonogenic progenitors was determined in aliquots of bone marrow cells as described above. After 10-14 days, individual colonies were plucked from the methylcellulose and processed to recover DNA. Specifically, after scoring the plates, 20 colonies (or less, if fewer were present) were picked at random and incubated in lysis buffer [50 nM Tris (pH 8.5), 1 mM EDTA, 0.5% Tween 20, and 100 ug/ml proteinase K] at 56°C for 2 hours. To inactivate the proteinase K, the samples were heated at 95°C for 10 minutes. The DNA samples were assayed for EGFP coding sequences using the polymerase chain reaction (PCR) methodology. PCR was performed with PCR Core Kit (Boehringer Mannheim) according to the manufacturer's instructions. The amplification conditions were as follows: 92°C for

2 min., then 35 cycles of 92°C for 1 min., 60°C for 1 min., and 72°C for 1 min., followed by a final elongation step of 7 min. at 72°C. Primers which amplify a 829 base pair (bp) fragment of human alpha satellite DNA, 5'-AATTCAGCTGACTAAACA-3' (SEQ ID NO:4) and 5'-TTTAGTTAGGTGCAGTTAT-3' (SEQ ID NO:2), were used to confirm the presence of human DNA in each sample. The PCR amplification of the EGFP gene was performed with the primers 5'-ACCCCGACCACATGAAGCAGC-3' (SEQ ID NO:3) and 5'-CGTTGGGGTCTTTGCTCAGGG-3' (SEQ ID NO:4), resulting in a 417 bp product. Primers specific to the β -actin gene, 5'-TGACGGGGTCACCCACACTGTGCCCATCTA-3' (SEQ ID NO:5) and 5'-CTAGAAGCATTTGCGGTGGACGATGGAGGG-3' (SEQ ID NO:6), were used as an internal control and gave a 604 bp product. The PCR products were electrophoresed on a 1% agarose gel with ethidium bromide staining to visualize DNA. Samples that failed to produce PCR fragments for either β -actin or alpha satellite DNA were not included in the calculation of gene transfer efficiency.

Results

Enhanced transduction of human hematopoietic stem cells with RD114 pseudotyped vector particles. The titer of infectious particles of various oncoviral vector preparations (*i.e.*, RD114, amphotropic, GALV, and VSV-G pseudotyped vector particles containing MGIRL22Y retroviral genome) was determined prior to transduction by limiting dilution on HeLa cells. Human cord blood CD34⁺ cells were cultured for 24 hours in cytokine-containing, serum-free medium (*see above*) and then transduced on retronectin coated plates. Serum-containing (10% FCS), conditioned medium from producer cells was added in amounts necessary to achieve the specified MOIs (but not exceeding 40% of the culture volume). After 24 hours fresh, serum-free cytokine-containing medium was added in amounts equal to the culture volume (2X dilution), and, after an additional 48 hours, the cells were analyzed for EGFP expression. With a single exposure at MOI of 5, the RD114 pseudotyped particles were far more efficient at transducing human CD34⁺ cells than were vector particles pseudotyped with the amphotropic, GALV, or VSV-G env proteins. Human T-lymphocytes activated by PHA and IL-2 and human leukemia cells (K562 and HEL) were also far more efficiently transduced with RD114 than with amphotropic pseudotyped particles.

Aliquots of cells transduced with either RD114 or amphotropic pseudotyped particles were replated in methylcellulose immediately following transduction (48 hours of culture) and incubated for an additional 10-15 days, and the resistance to trimetrexate was determined. The RD114 pseudotyped vector efficiently transduced CD34⁺ cells and their progenitors at a very low MOI indicating the presence of the appropriate receptor and cycling of a significant proportion of the cell population. Amphotropic pseudotyped particles failed to efficiently transduce the same population even at increased particle concentrations, consistent with a block to transduction at the receptor level.

Medium conditioned by derivatives of the HT1080 cell line alters the immunophenotype of primitive human hematopoietic cells during transduction. The immunophenotype of purified CD34⁺ and CD34⁺, CD38⁻ cells was routinely monitored following *in vitro* culture and transduction. After 96 hours in serum-free culture medium with high dose of cytokines, purified CD34⁺, CD38⁻ cells retained their phenotype (*see* also Bhatia *et al.*, J. Exp. Med., 1997, 186:619-624). In contrast, the exposure of CD34⁺, CD38⁻ cells to the medium conditioned by the RD114/MGirL22Y producer cells during transduction caused the majority of CD34⁺, CD38⁻ cells to become CD38⁺. While only a small decrease in total clonogenic progenitors accompanied this phenotypic change (p-NS), the inventors have found that CD34⁺ cells exposed to conditioned medium from the RD114/MGirL22Y producer cells failed to engraft in murine NOD/SCID recipients (N=4). In the same experiment, control CD34⁺ cells (cultured identically but without exposure to vector particles) exhibited multilineage engraftment.

The observed phenotypic changes during transduction and the loss of primitive, repopulating cells may be caused directly by the RD114 env protein or, alternatively, may arise due to the action of some other component within the medium conditioned by the RD114/MGirL22Y producer cells. The later possibility seems more likely in light of finding that the culture medium from an amphotropic producer cell population (derived from the same original cell line, HT1080, as the producer cells for RD114 pseudotyped particles; Sommerfelt *et al.*, Virology, 1990, 176:58-69), induced a similar immunophenotypic change in CD34⁺, CD38⁻ cells, *i.e.*, differentiation to a CD38⁺ phenotype. In contrast, the culture medium from an amphotropic producer cell line of a different origin (*i.e.*, derived from PA317 cells) did not affect the immunophenotype of the

CD34⁺, CD38⁻ cells. RD114 pseudotyped producer cells were also derived from 293T cells (*see* above). The culture medium from these cells had no effect on the immunophenotype of CD34⁺, CD38⁻ cells indicating that the immunodifferentiating/eliminating substance is indeed HT1080 cell-specific.

Transduction of primitive human hematopoietic cells by RD114

pseudotyped vector particles which had been pre-loaded onto retronectin coated plates.

To provide the proper conditions for engraftment of transduced hematopoietic stem cells in murine NOD/SCID recipients the inventors set to prevent the undesirable effect of HT1080 cell-specific substance inducing the immunophenotypic change of CD34⁺, CD38⁻ cells and the elimination of repopulating cells. Two alternative approaches were chosen.

According to the first approach, vector particles used for transduction were concentrated (and the volume of the culture medium containing harmful substances was greatly reduced) by ultracentrifugation at 25,000rpm for 90 minutes. Thus, in the experiment presented in the viral supernatant was concentrated 54-fold with highly efficient recovery of the particles (60%) without any loss of the transduction efficiency.

According to an alternative approach, retroviral vector particles were adsorbed or “pre-loaded” onto retronectin coated plates by a brief incubation with virus-conditioned medium (Williams, Ann. NY Acad. Sci., 1999, 872:109-113). In this way the vector particles become concentrated on the retronectin allowing the conditioned medium (containing a phenotype-altering substance) to be removed and replaced by medium containing the target cell population. Using conditioned medium from the RD114/MGirL22Y producer cells, the RD114 pseudotyped particles were adsorbed onto retronectin and then used to transduce CD34⁺, CD38⁻ cells. This experimental approach proved to eliminate the undesirable effect of the HT1080-derived culture medium as it maintained the immunophenotype of the CD34⁺, CD38⁻ cells with preservation of transduction efficiency. Transduction efficiency of pre-adsorbed viral particles remains unchanged after incubation for 48 hours at 4°C and upon long-term storage (greater than 1 week) at -70°C. Transduction efficiency of pre-adsorbed viral particles is cell concentration dependent with the optimal efficiency obtained at a concentration of 1-5 x 10⁴ cells/cm².

Further experiments using CD34⁺, CD38⁻ cells were performed to determine the optimum period of prestimulation prior to the transfer of the target cell

population to retronectin coated plates pre-loaded with RD114 pseudotyped particles (N=4). Significant transduction of clonogenic progenitors (as reflected by trimetrexate resistance) was achieved already after 24 hours of prestimulation ($38 \pm 15\%$). However, the maximal effect was seen only after 48 hours of prestimulation ($73 \pm 12\%$).

Transduction of human cells capable of establishing hematopoiesis in immunodeficient mice with RD114 pseudotyped vector particles. Purified CD34⁺ cells which had been prestimulated for 24-48 hours in serum-free medium were transduced by a single exposure to RD114 vector particles pre-loaded onto retronectin coated plates. After a maximum of 96 hours of culturing, expanded cells derived from an input $1.0-1.5 \times 10^5$ cells were injected into NOD/SCID immunodeficient murine recipients. Control cells were either not exposed to retroviral vector particles or transduced with amphotropic vector particles under various conditions. Bone marrow cells of three animals were analyzed 8-10 weeks after they had received CD34⁺ cells transduced with RD114 pseudotyped vector particles. In each animal, there was a substantial population of bone marrow cells which reacted with a human CD45-specific monoclonal antibody. Among these CD45⁺ cells, the EGFP-positive cells were found in both the human lymphoid (CD19⁺) and the human myeloid (CD33⁺) cell populations and in all three animals.

Overall rates of human engraftment in the NOD/SCID murine recipients and the levels of EGFP expression were determined. EGFP-positive cells were found in very significant numbers only in animals which received RD114 transduced CD34⁺ cells. The observed decrease in engraftment efficiency of CD34⁺ cells transduced by the RD114 pseudotyped vector after 24 hours of prestimulation (compared to untransduced CD34⁺ cells [$p < 0.05$]) was ameliorated by longer prestimulation (48 hours) which also resulted in improved EGFP expression. In correlation with reports of others (Glimm *et al.*, Blood, 1999, 94:2161-2168; Henneman *et al.*, Exp. Hematol., 1999, 27:817-825), a 48 hour-long prestimulation of producer cells prior to transduction appears to be an optimal condition for maximal gene transfer into NOD/SCID repopulating cells. The efficiency of gene transfer into repopulating hematopoietic cells was also evaluated by PCR analysis of DNA recovered from expanded bone marrow cells of experimental animals. A high percentage of cell clones from animals that had received cells transduced with RD114 pseudotyped vector particles produced EGFP-specific PCR product (Table 1). Of interest is the fact

to the fact that the same person may have been involved in more than one case.

Table 1

**Repopulation of NOD/SCID Mice by Transduced Cord Blood CD34⁺ Cells:
Analysis of Engraftment and Gene Transfer Efficiency**

Experimental Group	In Vitro Analysis		In Vivo Analysis			
	% EGFP Expression	% TMTX-R Progenitors	% Human CD45 ⁺ (range)	% EGFP ⁺ (range) ⁰	Human Progenitors	
					Trimetrexate Resistant (%)	PCR+ Colonies (%)
Expanded CD34⁺ Cells (N=10)⁺	0	0	13.5±17% (2.18-54.5%)	0%	0/228 (0%)	0/20 (0%)
AM/MGirL22Y Transduced CD34⁺ Cells (N=9)*	18-34%	6.5-8%	15.6±16% (1.0-53%)	0.2±0.1% (0-0.6%)	0/118 (0%)	0/50 (0%)
RD/MGirL22Y Preload Only @ 24 Hours (N=9)	75-87%	58-75%	3.5±5%^a (0.6-15.8%)	20.0±22%^b (0.6-71%)	24/106 (23%)	30/61 (49%)
RD/MGirL22Y Preload Only @ 48 Hours (N=6)	50-69%	45-66%	8.7±8% (1.7-23.5%)	40.5±40%^b (1.8-92%)	81/326 (25%)	58/98 (59%)

^a p<0.05 compared to controls recipients

^b p<0.05 compared to AM/MGirL22Y

⁰ Percent EGFP Expression of Engrafted Human Mononuclear Cells

⁺ N = Number of murine recipients of 1.0-1.5 x 10⁵ cells that exhibited > 0.5% human CD45⁺ cells

^{*} Transduction by preloading alone @ 24 hours (N=3) or with the addition of conditioned medium containing vector particles added at 48, and 72 hours

Discussion

The results presented above demonstrate that murine oncoretroviral vector particles pseudotyped with the envelope protein of feline endogenous virus (RD114) are

transduced into human primitive hematopoietic cell lines, the CD34⁺ and CD34⁺, CD38⁻ cell populations from cord blood, and the progenitors found in these populations far more efficiently than tested amphotropic, GALV and VSV-G pseudotyped oncoviral particles used at equivalent multiplicities of infection.

The disclosed producer cells generating the RD114 pseudotyped vector particles were derived from human sarcoma cells (HT1080) which apparently generate a substance that induces immunodifferentiation and depletion of repopulating cells. The nature and identity of this substance(s) is unknown and represents an important subject for future investigation. The evidence presented herein indicates that the RD114 pseudotyped particles are not the source of the substance since the same immunodifferentiation effect was observed with amphotropic particles produced by a derivative of the HT1080 cell line and was not observed with RD114 particles produced by human 293T cells.

As disclosed herein, the undesired effects of this substance could largely be avoided by preloading the RD114 particles onto retronectin plates prior to transduction of hematopoietic cell populations (although some depletion of NOD/SCID cells was observed compared to controls (Table 1). The ultimate solution may therefore involve the derivation of a RD114 packaging cell line from another cell type, such as NIH 3T3 mouse fibroblast cells which are currently the main source of vector preparations used in clinical trials.

As described above, cells in the cord blood CD34⁺ population transduced with retronectin-adsorbed RD114 pseudotyped vector particles are capable of establishing human hematopoiesis in immunodeficient (NOD/SCID) mice as reflected by the presence of the proviral genome in as many as 90% of the myeloid and lymphoid cells in the bone marrow of the transplant recipients. These results are far superior to gene transfer obtained to date with any onco- or lentiviral system.

The frequency of genetically modified, secondary progenitors present in bone marrow 8-10 weeks following transplantation was approximately equivalent to the frequency of genetically modified primary progenitors in the transduced population used for transplantation of the immunodeficient mice. These results imply that the frequency of transduction of NOD/SCID repopulating cells was equal to the transduction frequency of progenitors which, in turn, was equivalent to the transduction of the entire CD34⁺ cell population. In this respect, the results obtained in transducing human cord blood cells

with RD114 pseudotyped vector particles are comparable to those generally obtained in transducing murine hematopoietic cells with ecotropic vector particles (Allay *et al.*, *supra*; Persons *et al.*, *supra*). These results contrast with the general experience with amphotropic and GALV pseudotyped oncoretroviral vector particles in which the transduction frequency of more mature human CD34⁺ cells and progenitors exceeds that of repopulating cells (van Henn, *et al.*, Blood, 1998, 92:4013-4022; Marandin *et al.*, 1998, Hum. Gene Ther., 9:1497-1511; Schlitz *et al.*, Blood, 1998, 92:3163-3171; Conneally *et al.*, Blood, 1998, 91:3487-3493). Moreover, a higher frequency of transduction of human NOD/SCID repopulating cells was achieved with a single exposure to RD114 pseudotyped vector particles at a low MOI than with multiple exposures to amphotropic, GALV and VSV-G pseudotyped vector particles used at generally higher MOIs (*see also* van Henn, *et al.*, *supra*; Conneally *et al.*, *supra*; van Henn, *et al.*, *supra*; Marandin *et al.*, *supra*; Schlitz *et al.*, *supra*; Henneman *et al.*, *supra*; Rebel *et al.*, Blood, 1999, 93:2217-2224; Evans *et al.*, Hum. Gene Ther., 1999, 10:1479-1489). It can be therefore hypothesized that the neutral amino acid transporter which is used as a receptor for both RD114 and ecotropic pseudotyped particles (Albritton *et al.*, J. Virol., 1993, 67:2091-2096; Rasko *et al.*, Proc. Natl. Acad. Sci. USA, 1999, 96:2129-2134) is expressed at functionally higher levels on human primitive hematopoietic cells than is the phosphate transporter which serves as a receptor for amphotropic and GALV pseudotyped vector particles.

It has been reported that VSV-G pseudotyped lentiviral vector particles are superior to VSV-G pseudotyped oncoretroviral vector particles at transducing quiescent NOD/SCID repopulating cells (Miyoshi *et al.*, Science, 1999, 283:682-686). As discussed above, VSV-G pseudotyped oncoretroviral vector particles transduce NOD/SCID repopulating cells relatively inefficiently (Rebel *et al.*, *supra*) compared to the results obtained herein with RD114 pseudotyped vector particles. It can be inferred therefore that RD114 pseudotyped lentiviral vector particles may be even more superior than RD114 pseudotyped oncoviral vector particles at transducing primitive hematopoietic cells. Recent results from the inventors' laboratory suggest that the RD114 pseudotyped lentiviral vector particles can be generated.

Taken together, the disclosed data suggest that (i) increasing the transduction and repopulation frequency by pretreating the target cells with cytokines and by immobilizing the retroviral vector particles on retronectin plates, (ii) overcoming the limitation imposed

by receptor density for vector particle entry with the RD114-specific envelope protein, and (iii) the use of lentiviral vector internal proteins to facilitate translocation of the preintegration complex through the nuclear membrane may result in highly consistent and efficient transduction and repopulation of human hematopoietic stem cells thereby making gene therapy applications highly feasible and predictably successful.

EXAMPLE 2: Optimal Transduction Using RD114-Pseudotyped Vectors of Hematopoietic Stem Cells Requires Preloading/Adsorption of Vectors on Retronectin Coated Dishes and is Cell Concentration Dependent

As disclosed in Example 1 (*supra*), use of the RD114-pseudotyped retroviral vector particles pre-adsorbed (pre-loaded) onto retronectin-coated plates improves efficiency of stem cell transduction and NOD/SCID mice engraftment. It is further disclosed in Example 1 that the use of retronectin coated plates can eliminate the effects of unidentified producer cell-derived factors on the target stem cell differentiation and survival during engraftment.

In light of the above, it is important to attain the maximal efficiency of transduction when using retronectin-coated plates. We hypothesized that the transduction efficiency of the retronectin-adsorbed particles will be dependent on their density on the plate. Indeed, we found that gene transfer efficiency was maximized using retronectin concentration greater than 12 $\mu\text{g}/\text{cm}^2$. We further hypothesized that the transduction efficiency of the retronectin-adsorbed particles will be dependent on cell contact and, therefore, on cell concentration. Indeed, CD34⁺ cells from cord blood (CB) were efficiently transduced by adsorbed RD114-pseudotyped murine retroviruses on retronectin coated dishes at a concentration of $1-5 \times 10^4$ cells per cm^2 . Data were obtained from representative experiments of 2 cords. It has been repeated twice with peripheral blood CD34⁺ cells with similar results.

EXAMPLE 3: RD114-Pseudotyped Murine Retroviruses Adsorbed to Retronectin Coated Plates Maintain Transduction Efficiency of Hematopoietic Stem Cells After Long-Term Storage at -70°C/Dry Ice

We evaluated the stability of RD114-pseudotyped retroviral particles. We found that retroviruses pseudotyped with the RD114 envelope protein have a longer half-life in solution as compared to amphotropic pseudotyped retroviruses. We hypothesized that this stability would also be seen in the vector particles pre-loaded to the retronectin coated plates. We used frozen aliquots of a high titered RD114-MGirL22Y supernatant for these experiment. Peripheral blood CD34⁺ cells were transduced on 6 well plates prepared as follows: (i) pre-loaded and then frozen at 70°C for 48 hours before thawing, (ii) pre-loaded and placed at 4°C for 48hours, (iii) pre-loaded and placed at 4°C for 24 hours, or (iv) freshly pre-loaded with thawed supernatant. There was gradual loss of gene transfer efficiency using pre-loaded plates placed at 4°C as compared to pre-loaded plates prepared fresh. However, significant gene transfer efficiency could be maintained using pre-loaded plates that were kept frozen. We have subsequently repeated these experiments twice using plates stored at -70°C for greater than one week with similar results.

EXAMPLE 4: RD114-Pseudotyped Murine Retroviruses Can be Concentrated by Ultracentrifugation with Preservation of Transduction Efficiency

As disclosed in Example 1, to ensure the efficient retroviral-mediated gene transfer to stem cells, it is highly important to eliminate the effect of producer cell-derived factors on the target cell differentiation and survival during engraftment. Therefore, in parallel to developing the transduction protocols using viral particles pre-adsorbed on retronectin-coated plates (see Examples 1-3, *supra*), we separated the retroviral vector particles from the detrimental content of the producer cell culture media using ultracentrifugation.

The results show that RD114-pseudotyped vector particles can be collected in a concentrated form by ultracentrifugation at 25,000 rpm for 90 minutes at 4°C. In this experiment, the viral supernatant was concentrated 54-fold with 60% recovery of total particles.

The concentrated RD114 supernatant demonstrated greatly increased transduction efficiency when tested on HeLa cells. AZT was added to cells as a control to rule out pseudo-transduction. To determine the transduction efficiency, double color Facs

analysis was performed using peripheral blood transduced at 48 hours with GFP-pre-loaded virus, GFP-pre-loaded virus plus concentrated yellow fluorescent protein (YFP) plus supernatant at 48 hours, GFP-pre-loaded virus at 48 hours plus concentrated YFP plus supernatant at 72 hours, and GFP-pre-loaded virus at 72 hours.

5

EXAMPLE 5: RD114 Envelope Protein Pseudotyped Lentiviral Constructs Can Be Concentrated by Ultracentrifugation

As discussed in Example 1 (*supra*), the development of lentiviral-based retroviral vector particles can greatly improve the efficiency of gene transfer to the non-
10 dividing cells. We have therefore set to obtain RD114-pseudotyped lentiviral vector particles which can be used for transduction of various quiescent stem cells.

We have first prepared lentiviral vectors transiently using VSV-G pseudotyping and have observed, as have others (*e.g., see* Evans, *et al.*, Hum. Gene Ther.,
15 1999, 10:1479-1489), that these vectors can efficiently transduce CD34⁺,CD38⁻ cells while oncoretroviral particles pseudotyped with VSV-G do not.

We then sought to determine whether the RD114 envelope can pseudotype lentiviruses. We used two different transient transfection protocols and simultaneously derived VSV-G pseudotyped lentiviral vectors as an internal control to show that the
20 transfection protocols worked. Supernatants from producer cells were collected and viral titers were determined using HeLa cells. We found that we could indeed pseudotype lentiviruses with the RD114 envelope protein.

While in the first series of experiments the titer of the RD114 pseudotyped lentiviral preparation was low, viral transduction did occur. We have determined that the
25 low titer was the result of the poor expression of the envelope protein in the chosen producer system. In light of this observation, we have subsequently derived RD114-pseudotyped lentiviral particles with a much higher titer using a stronger promoter to drive the expression of RD114 envelope protein.

In addition, we have demonstrated that the RD114-pseudotyped lentiviral
30 vector particles can be concentrated (as described in Examples 1-4, *supra*) by pre-loading on retronectin-coated plates or by ultracentrifugation.

EXAMPLE 6: Repopulation of Hematopoietic Lineages of Immunodeficient Rhesus Monkey by Cord Blood CD34⁺ Cells Transduced with RD114-pseudotyped Retroviral Vector Particles Pre-adsorbed on Retronectin-coated Plates

5

Materials and Methods

G-CSF/SCF mobilized rhesus monkey peripheral blood CD34⁺ cells were selected and purified as described by Donahue *et al.* (Blood, 2000, 95:445-452). These cells were pre-stimulated under serum-free conditions essentially as described in Example 1 for NOD/SCID-repopulating cells (*supra*), except that the cytokines were SCF (300ng/ml), Flt-3 ligand (300 ng/ml), and interleukin-6 (50ng/ml). Following 48 hour pre-stimulation, cells were placed on retronectin-coated plates pre-loaded with RD-114-pseudotyped retroviral vector particles (RD114-MGirL22Y; prepared and pre-loaded as described in Examples 1 and 2, respectively) along with fresh media and cytokines. Upon overnight incubation (72 hours total), cells were removed from the plates, pelleted by centrifugation, and resuspended again in fresh media supplemented with cytokines. Cell suspension was plated for the second time on fresh retronectin-coated plates pre-loaded with retroviral vector particles. Cells were harvested at 96 hours, washed with fresh media, and injected into immunodeficient rhesus monkeys.

20

Results

The efficiency of engraftment of CD34⁺ cells transduced with RD114 pseudotyped vector particles in the hematopoietic lineages of the recipient monkey (96E113) was analyzed 0-106 days post-injection by monitoring EGFP expression. The percentage of hematopoietic cells expressing EGFP reached the maximal levels 64 days after injection, with the highest frequency of engraftment observed in monocytes (up to 98%) and peripheral mononuclear lymphocytes (PMN) (up to 83%). Lymphocytes and platelets also showed increased levels of EGFP expression, although not at the same level as monocytes or PMNs. Also, platelets achieved maximal EGFP expression at day 78 post transplantation.

30

* * *

The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described
5 herein will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims.

It is further to be understood that all values are approximate, and are provided for description.

All patents, applications, publications, test methods, literature, and other
10 materials cited herein are hereby incorporated by reference.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2